

XP-002543705

C:\EPOPROGS\SEA\1.1.1.epodata\sealeplogflinternal.log

## WPI / Thomson

AN - 1983-26776K [11]  
 AP - JP19730121394 19731029  
 CPY - MATS-I  
 DC - B04  
 DW - 198311  
 IC - A61K35/78  
 LNKA- 1983-026238  
 M1 - [01] M423 M781 M903 P816 V400 V404  
 MC - B04-A07D B12-H05  
 PA - (MATS-I) MATSUI T  
 PH - JP58009084B B 19830218 DW198311  
 JP50070512 A 19750612 DW198311  
 PR - JP19730121394 19731029  
 XIC - A61K-035/78; A61K-036/185; A61K-036/60; A61P-003/00; A61P-003/08;  
 C12C-003/00; C12C-007/00  
 AB - Cones of hop obtd. within 35-45 days from flowering are dried  
 naturally without exposure to sun or by exposing to air at 40-60 deg.C  
 for 5-15 hrs., adjusting moisture content to 10-20%. The cones are  
 crushed after drying, and the prod. used in the treatment of diabetes.  
 (J50070512)  
 ICAI- A61K36/60; A61P3/08; C12C3/00; C12C7/00  
 ICCI- A61K36/185; A61P3/00; C12C3/00; C12C7/00  
 IW - POWDER HOP EXTRACT PREPARATION TREAT DIABETES  
 IWW - POWDER HOP EXTRACT PREPARATION TREAT DIABETES  
 NC - 1  
 NPH - 2  
 OPD - 1973-10-29  
 PAW - (MATS-I) MATSUI T  
 PD - 1975-06-12  
 TI - Powdered hop extract prepn. - used for treatment of diabetes (J5  
 12.6.75)